

# A short proof of the tree-packing theorem

Tomáš Kaiser\*

## Abstract

We give a short elementary proof of Tutte and Nash-Williams' characterization of graphs with  $k$  edge-disjoint spanning trees.

We deal with graphs that may have parallel edges and loops; the vertex and edge sets of a graph  $H$  are denoted by  $V(H)$  and  $E(H)$ , respectively. Let  $G$  be a graph. If  $\mathcal{P}$  is a partition of  $V(G)$ , we let  $G/\mathcal{P}$  be the graph on the set  $\mathcal{P}$  with an edge joining distinct vertices  $P_1, P_2 \in \mathcal{P}$  for every edge of  $G$  with one end in  $P_1$  and another in  $P_2$ . Tutte [5] and Nash-Williams [3] proved the following classical result:

**Theorem 1.** *A graph  $G$  contains  $k$  pairwise edge-disjoint spanning trees if and only if for every partition  $\mathcal{P}$  of  $V(G)$ , the graph  $G/\mathcal{P}$  has at least  $k(|\mathcal{P}| - 1)$  edges.*

An elegant proof of Theorem 1 is based on the matroid union theorem (see, e.g., [4, Corollary 51.1a]); a relatively short elementary proof appears in [1, Theorem 2.4.1]. In this paper, we give another elementary proof which is also short and perhaps somewhat more straightforward. The argument directly translates to an efficient algorithm to find either  $k$  disjoint spanning trees, or a proof that none exist. To an extent, the method can also be applied to the packing of structures without the matroidal properties of spanning trees, as shown, e.g., in the forthcoming paper [2].

Let  $k \geq 1$  and  $\mathcal{T} = (T_1, \dots, T_k)$  be an ordered partition of  $G$  into  $k$  spanning subgraphs of  $G$ . We define the sequence  $\mathcal{P}_0, \mathcal{P}_1, \dots, \mathcal{P}_\infty$  of partitions of  $V(G)$  associated with  $\mathcal{T}$  as follows. First,  $\mathcal{P}_0 = \{V(G)\}$ . For  $i \geq 0$ , if there is  $c \in \{1, \dots, k\}$  such that the induced subgraph  $T_c[P]$  is disconnected for some  $P \in \mathcal{P}_i$ , then  $c_i$  is defined as the least such  $c$  and  $\mathcal{P}_{i+1}$  consists of the vertex sets of all

---

\*Department of Mathematics and Institute for Theoretical Computer Science (ITI), University of West Bohemia, Univerzitní 8, 306 14 Plzeň, Czech Republic. E-mail: kaisert@kma.zcu.cz. Supported by projects 1M0545, ME 885 and Research Plan MSM 4977751301 of the Czech Ministry of Education.

components of  $T_{c_i}[Q]$ , where  $Q$  ranges over all the classes of  $\mathcal{P}_i$ . Otherwise, the process ends by setting  $\mathcal{P}_\infty = \mathcal{P}_i$ .

We define the *level*  $\ell(e)$  of an edge  $e \in E(G)$  (with respect to  $\mathcal{T}$ ) as the largest  $i$  (possibly  $\infty$ ) such that both ends of  $e$  are contained in one class of  $\mathcal{P}_i$ . An edge  $e \in E(T_k)$  is *superfluous* (for  $\mathcal{T}$ ) if  $\ell(e) < \infty$  and  $e$  is contained in some cycle of  $T_k$ . We say that an edge *leaves*  $P \subset V(G)$  if it has precisely one end in  $P$ .

To keep the notation simple, the symbols  $\mathcal{P}_i$  and  $\ell(e)$  (as well as  $\mathcal{P}_\infty$  and  $c_i$ ) will relate to a partition  $\mathcal{T}$ , while  $\mathcal{P}'_i$  and  $\ell'(e)$  relate to a partition  $\mathcal{T}'$ . Thus, for instance, the level  $\ell'(e)$  of an edge  $e$  with respect to  $\mathcal{T}'$  is defined using the partitions  $\mathcal{P}'_i$  associated with  $\mathcal{T}'$ .

*Proof of Theorem 1.* The necessity of the condition is clear. To prove the sufficiency, we proceed by induction on  $k$ . The claim is trivially true for  $k = 0$ , so assume  $k \geq 1$  and choose an ordered partition  $\mathcal{T} = (T_1, \dots, T_k)$  of  $G$  into spanning subgraphs of  $G$  such that  $T_1, \dots, T_{k-1}$  are trees and the following holds:

- (1) the number of components of  $T_k$  is as small as possible,
- (2) either there is no superfluous edge for  $\mathcal{T}$ , or the minimum level of a superfluous edge is as small as possible subject to (1).

If  $T_k$  is connected, then we are done. For the sake of a contradiction, suppose that  $T_k$  has at least two components (i.e.,  $|\mathcal{P}_1| \geq 2$ ). We prove that there exists a superfluous edge for  $\mathcal{T}$ . For all  $i = 1, \dots, k-1$  and  $P \in \mathcal{P} := \mathcal{P}_\infty$ , the graph  $T_i[P]$  is connected. Hence  $T_i/\mathcal{P}$  is a tree and has exactly  $|\mathcal{P}| - 1$  edges. By the assumption on  $G$ , the graph  $T_k/\mathcal{P}$  has at least  $k(|\mathcal{P}| - 1) - (k-1)(|\mathcal{P}| - 1) = |\mathcal{P}| - 1$  edges. Since  $T_k/\mathcal{P}$  has  $|\mathcal{P}|$  vertices and is disconnected, it must contain a cycle. But then  $T_k$  contains a cycle as each  $T_k[P]$  is connected for  $P \in \mathcal{P}$ . All the edges of this cycle joining different classes of  $\mathcal{P}$  (there are at least two such edges) are superfluous for  $\mathcal{T}$ .

Let  $e \in E(T_k)$  be an edge of minimum level that is superfluous for  $\mathcal{T}$  and set  $m = \ell(e)$ . Let  $P$  be the class of  $\mathcal{P}_m$  containing both ends of  $e$ . Since  $e$  joins different components of  $T_{c_m}[P]$ , we have  $c_m \neq k$  and the unique cycle  $C$  in  $T_{c_m} + e$  contains an edge leaving  $P$ . Thus, for an edge  $e'$  of  $C$  of lowest possible level we have  $j := \ell(e') < \ell(e) = m$ . Let  $Q$  be the class of  $\mathcal{P}_j$  containing both ends of  $e'$ ; note that (the vertex set of)  $C$  is contained within  $Q$ . We will exchange  $e$  for  $e'$  in the respective sets  $T_i$  to eventually obtain the desired contradiction.

Let  $\mathcal{T}'$  be the partition obtained from  $\mathcal{T}$  by replacing  $T_{c_m}$  with  $T'_{c_m} = T_{c_m} + e - e'$  and  $T_k$  with  $T'_k = T_k - e + e'$ ; we set  $T'_i = T_i$  for  $i \notin \{c_m, k\}$ . If the ends of  $e'$  are in different components of  $T_k$  (i.e.,  $j = 0$ ), then  $T_k - e + e'$  has fewer components than  $T_k$ , which is impossible. Thus, all of  $C$  is contained within one class  $R \in \mathcal{P}_1$ . Furthermore, since  $e$  is superfluous for  $\mathcal{T}$ ,  $T_k[R] - e$  is connected and so  $e'$  is superfluous for  $\mathcal{T}'$ .

**Claim.** For all  $i \leq j$ ,  $\mathcal{P}'_i = \mathcal{P}_i$ .

The proof is by induction on  $i$ , with the  $i = 0$  case being trivial. We assume that  $\mathcal{P}'_i = \mathcal{P}_i$  and prove that every component of  $T_c[U]$ , where  $U \in \mathcal{P}_i$  and  $c \leq c_i$ , has the same vertex set as some component of  $T'_c[U]$ . This will clearly imply the Claim.

Let  $K$  be the vertex set of a component of  $T_c[U]$ . First of all,  $T'_c$  contains no edge leaving  $K$ , because  $T_c$  and  $T'_c$  only differ by edges contained within  $Q \in \mathcal{P}_j$ , which is either disjoint from  $K$  or contained in it. Thus, it suffices to prove that  $T'_c[K]$  is connected. Furthermore, we may restrict our attention to  $c \in \{c_m, k\}$ . Suppose that  $T'_c[K]$  is disconnected; there are two cases.

Case 1:  $c = k$  and  $T_k[K] - e$  is disconnected. Since  $e$  is superfluous, it is contained in a cycle  $D$  of  $T_k$ . By the assumption,  $D$  is not contained within  $K \in \mathcal{P}_i$ , so some edge  $f$  of  $D$  leaves  $K$ . However,  $f$  is then superfluous for  $\mathcal{T}$  and  $\ell(f) \leq i < m$ , a contradiction.

Case 2:  $c = c_m$  and  $T_{c_m}[K] + e - e'$  is disconnected. The class  $Q \in \mathcal{P}_j$  containing the cycle  $C$  is thus contained in  $K$ . But then  $C$  is a cycle in  $T_{c_m}[K] + e$  containing  $e'$ , so  $T_{c_m}[K] + e - e'$  cannot be disconnected. This contradiction concludes the proof of the Claim.

We continue with the proof of the theorem. The above Claim implies that  $\ell'(e') \geq j$ . On the other hand, if  $c_j \neq k$ , then  $e'$  joins different components of  $T'_{c_j}[Q]$  and so we actually have  $\ell'(e') = j < i$ , a contradiction with the choice of  $e$  since we have seen that  $e'$  is superfluous for  $\mathcal{T}'$ . Thus, we conclude that  $c_j = k$ . Since  $T_k[R]$  is connected, the ends of  $e'$  are joined by a path  $S$  in  $T_k[R]$ . However,  $S$  is not contained within  $Q$  as  $e'$  joins different components of  $T_k[Q]$ . Hence,  $S$  contains an edge  $e''$  leaving  $Q$ . Observing that  $\ell(e'') < j$ , we use the Claim to obtain  $\ell'(e'') = \ell(e'') < j < \ell(e)$ , a final contradiction with the minimality of  $e$  since  $e''$  is superfluous for  $\mathcal{T}'$ .  $\square$

## References

- [1] R. Diestel, *Graph Theory*, 3rd Edition, Springer, 2005.
- [2] T. Kaiser and P. Vrána, Hamilton cycles in 6-connected line graphs, in preparation.
- [3] C. St. J. A. Nash-Williams, Edge-disjoint spanning trees of finite graphs, *J. London Math. Soc.* **36** (1961), 445–450.
- [4] A. Schrijver, *Combinatorial Optimization*, Springer, 2003.
- [5] W. T. Tutte, On the problem of decomposing a graph into  $n$  connected factors, *J. London Math. Soc.* **36** (1961), 221–230.