

Characterizing forbidden disjoint pairs for 2-factor of 2-connected graphs

Shipeng Wang

Beijing Institute of Technology

Beijing, China

12th April, Plzen, Czech

Joint work with Přemysl Holub, Zdeněk Ryjáček, Petr Vrána and

Liming Xiong

Terminology

All graphs are finite, simple and undirected.

$\alpha(G)$: the independence number of G ,

$\kappa(G)$: the connectivity of G ,

$G[X]$: the subgraph of G induced by X ,

$N_G(x)$: the neighborhood of x in G ,

$N_C(S)$: the neighborhood of S in the subgraph C of G .

Terminology

- A graph G is **hamiltonian**: G has a spanning cycle.
- A **2-factor** of a graph is a spanning subgraph whose components are cycles.
- The **Ramsey number** $r(k, l)$ is defined as the smallest integer such that every graph on $r(k, l)$ vertices contains either a clique of k vertices or an independent set of l vertices.

Terminology

- Let \mathcal{H} be a set of graphs. A graph G is said to be \mathcal{H} -free if G does not contain F as an induced subgraph for all F in \mathcal{H} , and we call each graph F of \mathcal{H} a **forbidden subgraph**.
- Let \mathcal{H} be a set of two graphs. Then we call \mathcal{H} a **forbidden disjoint pair**. Specially, if each of \mathcal{H} is connected, then we call \mathcal{H} a **forbidden pair**.

Terminology

- We use $H_1 \cup H_2$ to denote the **disjoint union** of two vertex-disjoint graphs H_1 and H_2 .
- We use $H_1 \cup H_2$ -free instead of $(H_1 \cup H_2)$ -free.
- So, $H_1 \cup H_2$ -free means we forb $H_1 \cup H_2$ as an induced subgraph; it does not mean forbidding H_1 and/or H_2 .

Motivation

Theorem 1.1 (Bedrossian, 1991)

Let $\{R, S\}$ be a forbidden pair such that neither R nor S is an induced subgraph of P_3 . Then every 2-connected graph \mathcal{H} -free graph G is hamiltonian if and only if (up to symmetry) $R = K_{1,3}$ and S is an induced subgraph of P_6 , $N_{1,1,1}$, or $B_{1,2}$.

a

^aBedrossian, Forbidden subgraph and minimum degree conditions for Hamiltonicity, Ph.D. Thesis, Memphis State University, 1991




Figure: Z_n , $B_{m,n}$ and $N_{l,m,n}$

Motivation

Because some non-Hamiltonian 2-connected graph with **small order**, Faudree and Gould have characterized all forbidden pairs for **hamiltonicity** of 2-connected graphs with **sufficiently large order**.

Theorem 1.2 (Faudree and Gould, DM, 1997)

Let $\{R, S\}$ be a forbidden pair such that neither R nor S is an induced subgraph of P_3 . Then every 2-connected graph \mathcal{H} -free graph with sufficiently large order is hamiltonian if and only if (up to symmetry) $R = K_{1,3}$ and S is an induced subgraph of $P_6, N_{1,1,1}, B_{1,2}$ or Z_3 .

^a

^aR. J. Faudree and R. J. Gould, Characterizing forbidden pairs for Hamiltonian properties, Discrete Math. 173(1997)45-60

Motivation

Because some non-Hamiltonian 2-connected graph with **small order**, Faudree and Gould have characterized all forbidden pairs for **hamiltonicity** of 2-connected graphs with **sufficiently large order**.

Theorem 1.2 (Faudree and Gould, DM, 1997)

Let $\{R, S\}$ be a forbidden pair such that neither R nor S is an induced subgraph of P_3 . Then every 2-connected graph \mathcal{H} -free graph with sufficiently large order is hamiltonian if and only if (up to symmetry) $R = K_{1,3}$ and S is an induced subgraph of $P_6, N_{1,1,1}, B_{1,2}$ or Z_3 .

^a

^aR. J. Faudree and R. J. Gould, Characterizing forbidden pairs for Hamiltonian properties, Discrete Math. 173(1997)45-60

Motivation

Faudree, Faudree and Ryjáček characterized all forbidden pairs for **2-factor** of 2-connected graphs with **sufficiently larger order**.

Theorem 1.3 (Faudree, Faudree and Ryjáček, DM, 2008)

Let $\{R, S\}$ be a forbidden pair such that neither R nor S is an induced subgraph of P_3 . Then every 2-connected graph $\{R, S\}$ -free graph with sufficiently large order has a 2-factor (up to symmetry) $R = K_{1,3}$ and S is an induced subgraph of $P_7, B_{1,4}, N_{1,1,3}$, or $R = K_{1,4}$ and $S = P_4$.

^a

^aJ. R. Faudree, R. J. Faudree and Z. Ryjáček, Forbidden subgraphs that imply 2-factors, Discrete Math. 308(2008)1571-1582.

Motivation

Li and Vrána characterized all forbidden disjoint pairs for **hamiltonicity** of 2-connected graphs.

Theorem 1.4 (Li and Vrána, JGT, 2017)

Let $\{R, S\}$ be a forbidden disjoint pair such that neither R nor S is an induced subgraph of P_3 or $3K_1$. Then there is an integer n_0 such that every 2-connected $\{R, S\}$ -free graph of order at least n_0 is hamiltonian, if and only if, one of the following is true (up to symmetry):

- $R = K_{1,3}$ and S is an induced subgraph of P_6 , Z_3 , $N_{0,1,2}$, $N_{1,1,1}$, $K_1 \cup Z_2$, $K_2 \cup Z_1$, or $K_3 \cup P_4$;
- $R = K_{1,k}$ with $k \geq 4$ and S is an induced subgraph of $2K_1 \cup K_2$;
- $R = kK_1$ with $k \geq 4$ and S is an induced subgraph of L_l with $l \geq 3$, or $2K_1 \cup K_{l'}$ with $l' \geq 2$.

Our results

We consider the following problem.

Problem: Determine the set \mathcal{H} of forbidden disjoint pairs \mathcal{H} such that every 2-connected \mathcal{H} -free graph G with sufficiently larger order has a 2-factor.

We first characterized forbidden subgraphs to guarantee a 2-connected graph with sufficiently large order has a 2-factor.

Theorem 2.1 (Holub, Ryjáček, Vrána, Wang and Xiong, 2017⁺)

Let H be a graph. Then every 2-connected H -free graph G with order at least $r(4, 28)$ has a 2-factor if and only if H is an induced subgraph of P_3 or $4K_1$.

Our results

We consider the following problem.

Problem: Determine the set \mathcal{H} of forbidden disjoint pairs \mathcal{H} such that every 2-connected \mathcal{H} -free graph G with sufficiently larger order has a 2-factor.

We first characterized forbidden subgraphs to guarantee a 2-connected graph with sufficiently large order has a 2-factor.

Theorem 2.1 (Holub, Ryjáček, Vrána, Wang and Xiong, 2017⁺)

Let H be a graph. Then every 2-connected H -free graph G with order at least $r(4, 28)$ has a 2-factor if and only if H is an induced subgraph of P_3 or $4K_1$.

Our results

Theorem 2.2 (Holub, Ryjáček, Vrána, Wang and Xiong, 2017⁺)

Let $\{R, S\}$ be a forbidden disjoint pair such that neither R nor S is an induced subgraph of P_3 or $4K_1$. Then there is an integer n_0 such that every 2-connected $\{R, S\}$ -free graph of order at least n_0 has a 2-factor if and only if (up to symmetry):

- $R = K_{1,3}$ and S is an induced subgraph of P_7 , $B_{1,4}$, $N_{1,1,3}$, $\Delta \cup Z_1$, $Z_1 \cup P_4$, or $Z_4 \cup K_1$;
- $R = K_{1,4}$ and S is an induced subgraph of $P_3 \cup 2K_1$;
- $R = K_{1,r}$ with $r \geq 5$ and S is an induced subgraph of $P_3 \cup K_1$ or $3K_1 \cup K_2$;
- $R = kK_1$ with $k \geq 5$ and S is an induced subgraph of L_l with $l \geq 3$, or $3K_1 \cup K_{l'}$ with $l' \geq 2$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Next, we will sketch the proof of one pair $\{K_{1,4}, P_3 \cup 2K_1\}$. Before sketching the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$, we need the following theorems and lemma.

Theorem 2.3 (Chvátal and Erdős, 1972, DM)

Let G be a graph on at least three vertices with independent number α and connectivity κ , where $\alpha \leq \kappa$. Then G is hamiltonian.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Next, we will sketch the proof of one pair $\{K_{1,4}, P_3 \cup 2K_1\}$. Before sketching the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$, we need the following theorems and lemma.

Theorem 2.3 (Chvátal and Erdős, 1972, DM)

Let G be a graph on at least three vertices with independent number α and connectivity κ , where $\alpha \leq \kappa$. Then G is hamiltonian.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Theorem 2.4 (Aldred, Egawa, Fujisawa, Ota and Saito, 2011, JGT)

For $n \geq 4$, every $(n-1)$ -connected $K_{1,n}$ -free graph contains a 2-factor.

Lemma 2.5 (Holub, Ryjáček, Vrána, Wang and Xiong, 2017⁺)

Let G be a 2-connected kK_1 -free graph such that G can be partitioned into two sets X and Y satisfying the following:

- (1) X contains a clique C such that each vertex of X has at least $k+6$ neighbors in C ;
- (2) $\alpha(G[Y]) \leq 2$.

Then G has a 2-factor.

Now, we will prove that every 2-connected $\{K_{1,4}, P_3 \cup 2K_1\}$ -free graph with order at least $r(5, r(4, 28) + 4)$ has a 2-factor.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Theorem 2.4 (Aldred, Egawa, Fujisawa, Ota and Saito, 2011, JGT)

For $n \geq 4$, every $(n - 1)$ -connected $K_{1,n}$ -free graph contains a 2-factor.

Lemma 2.5 (Holub, Ryjáček, Vrána, Wang and Xiong, 2017⁺)

Let G be a 2-connected kK_1 -free graph such that G can be partitioned into two sets X and Y satisfying the following:


- (1) X contains a clique C such that each vertex of X has at least $k + 6$ neighbors in C ;
- (2) $\alpha(G[Y]) \leq 2$.

Then G has a 2-factor.

Now, we will prove that every 2-connected $\{K_{1,4}, P_3 \cup 2K_1\}$ -free graph with order at least $r(5, r(4, 28) + 4)$ has a 2-factor.


Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- We first claim that G is $5K_1$ -free.
- Since G is $5K_1$ -free and $n(G) \geq r(5, r(4, 28) + 4)$, G contains a clique C of size $r(4, 28) + 4$. Set $X = \{x \in V(G) : d_C(x) \geq 16\}$ and $Y = V(G) \setminus X$, clearly $C \subseteq X$. Then $\kappa(G[X]) \geq 16$ and $\alpha(G[X]) \leq \alpha(G) \leq 4$, $G[X]$ is hamiltonian by Chvátal-Erdős Theorem.
- We can prove that $\alpha(G[Y]) \leq 3$. If $\alpha(G[Y]) \leq 2$, then by Lemma 2.5, we are done. Hence we assume that $\alpha(G[Y]) = 3$.



Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- We first claim that G is $5K_1$ -free.
- Since G is $5K_1$ -free and $n(G) \geq r(5, r(4, 28) + 4)$, G contains a clique C of size $r(4, 28) + 4$. Set $X = \{x \in V(G) : d_C(x) \geq 16\}$ and $Y = V(G) \setminus X$, clearly $C \subseteq X$. Then $\kappa(G[X]) \geq 16$ and $\alpha(G[X]) \leq \alpha(G) \leq 4$, $G[X]$ is hamiltonian by Chvátal-Erdős Theorem.
- We can prove that $\alpha(G[Y]) \leq 3$. If $\alpha(G[Y]) \leq 2$, then by Lemma 2.5, we are done. Hence we assume that $\alpha(G[Y]) = 3$.



Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

In the following, we need distinguish two cases.

Case 1. $G[Y]$ is disconnected.

By $\alpha(G[Y]) = 3$, $G[Y]$ has at most three components. We can prove that $G[Y]$ has exactly three components Y_1, Y_2, Y_3 and each $G[Y_i]$ is a clique.

For $i \in \{1, 2, 3\}$, let $P(z_1^i, z_2^i) = z_1^i P^i z_2^i$ denote a path of $G[X \cup Y_i]$ such that $V(P^i) = V(Y_i)$ and $z_1^i, z_2^i \in X$, and let $X_1 = V(P(z_1^1, z_2^1) \cap P(z_1^2, z_2^2) \cap P(z_1^3, z_2^3))$.

We need to consider the following six basic cases because any other cases can be changed to some basic case.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

In the following, we need distinguish two cases.

Case 1. $G[Y]$ is disconnected.


By $\alpha(G[Y]) = 3$, $G[Y]$ has at most three components. We can prove that $G[Y]$ has exactly three components Y_1, Y_2, Y_3 and each $G[Y_i]$ is a clique.

For $i \in \{1, 2, 3\}$, let $P(z_1^i, z_2^i) = z_1^i P^i z_2^i$ denote a path of $G[X \cup Y_i]$ such that $V(P^i) = V(Y_i)$ and $z_1^i, z_2^i \in X$, and let $X_1 = V(P(z_1^1, z_2^1) \cap P(z_1^2, z_2^2) \cap P(z_1^3, z_2^3))$.

We need to consider the following six basic cases because any other cases can be changed to some basic case.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$


Subcase 1.1. $|X_1| = 6$.



Since $\kappa(G[X - X_1 \cup X_2]) \geq \kappa(G[X]) - |X_1 \cup X_2| \geq 16 - 12 = 4$ and $\alpha(G[X - X_1 \cup X_2]) \leq \alpha(G) \leq 4$, $G[X - X_1 \cup X_2]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly four components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$



Subcase 1.2. $|X_1| = 5$.



Since $\kappa(G[X - X_1 \cup X_2]) \geq \kappa(G[X]) - |X_1 \cup X_2| \geq 16 - 9 = 7$ and $\alpha(G[X - X_1 \cup X_2]) \leq \alpha(G) \leq 4$, $G[X - X_1 \cup X_2]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly three components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$


Subcase 1.3. $|X_1| = 4$.



Since $\kappa(G[X - X_1 \cup X_2]) \geq \kappa(G[X]) - |X_1 \cup X_2| \geq 16 - 9 = 7$ and $\alpha(G[X - X_1 \cup X_2]) \leq \alpha(G) \leq 4$, $G[X - X_1 \cup X_2]$ is hamiltonian by Chvátal-Erdős Theorem, and thus G has a 2-factor.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 1.4. $|X_1| = 3$.



Since $\kappa(G[X - X_1]) \geq \kappa(G[X]) - |X_1| \geq 16 - 3 = 13$ and $\alpha(G[X - X_1]) \leq \alpha(G) \leq 4$, $G[X - X_1]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly two components.


Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Case 2. $G[Y]$ is connected.

If $\kappa(G[Y]) \geq 3$, then $G[Y]$ is 3-connected $K_{1,4}$ -free, $G[Y]$ has a 2-factor by Theorem 2.4, and thus G has a 2-factor.

Subcase 2.1. $\kappa(G[Y]) = 2$.

Let $\{v_1, v_2\}$ be a vertex-cut of $G[Y]$. By $\alpha(G[Y]) = 3$, $G[Y] - \{v_1, v_2\}$ has at most three components. If $G[Y] - \{v_1, v_2\}$ has exactly two components Y_1, Y_2 , then we can prove that both $G[Y_1]$ and $G[Y_2]$ are cliques. Hence $G[Y]$ is hamiltonian, and thus G has a 2-factor with exactly two components.




Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Case 2. $G[Y]$ is connected.

If $\kappa(G[Y]) \geq 3$, then $G[Y]$ is 3-connected $K_{1,4}$ -free, $G[Y]$ has a 2-factor by Theorem 2.4, and thus G has a 2-factor.

Subcase 2.1. $\kappa(G[Y]) = 2$.


Let $\{v_1, v_2\}$ be a vertex-cut of $G[Y]$. By $\alpha(G[Y]) = 3$, $G[Y] - \{v_1, v_2\}$ has at most three components. If $G[Y] - \{v_1, v_2\}$ has exactly two components Y_1, Y_2 , then we can prove that both $G[Y_1]$ and $G[Y_2]$ are cliques. Hence $G[Y]$ is hamiltonian, and thus G has a 2-factor with exactly two components.



Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Hence we assume that $G[Y] - \{v_1, v_2\}$ has exactly three components Y_1, Y_2, Y_3 . Then by $\alpha(G[Y]) = 3$, each $G[Y_i]$ is a clique.

- If some Y_i has at least three vertices, then $G[Y_i]$ has hamiltonian cycle. Thus we delete the set Y_i and the resulting graph of $G[Y] - Y_i$ has a hamiltonian cycle, and thus G has a 2-factor with exactly three components.




- Hence we assume that $|Y_i| \leq 2$ for $i \in \{1, 2, 3\}$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Hence we assume that $G[Y] - \{v_1, v_2\}$ has exactly three components Y_1, Y_2, Y_3 . Then by $\alpha(G[Y]) = 3$, each $G[Y_i]$ is a clique.

- If some Y_i has at least three vertices, then $G[Y_i]$ has hamiltonian cycle. Thus we delete the set Y_i and the resulting graph of $G[Y] - Y_i$ has a hamiltonian cycle, and thus G has a 2-factor with exactly three components.



- Hence we assume that $|Y_i| \leq 2$ for $i \in \{1, 2, 3\}$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- We claim that v_i is adjacent to every vertex in $Y_1 \cup Y_2 \cup Y_3$ for $i \in \{1, 2\}$. Since G is 2-connected, there exists two disjoint edges x_1y_1, x_2y_2 such that $x_1, x_2 \in X$ and $y_1, y_2 \in \{v_1, v_2\} \cup Y_1 \cup Y_2$.

We need to consider the following three basic cases because any other cases can be changed to some basic case.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 2.1.1. y_1 and y_2 are in the same Y_i .




Figure: G has a 2-factor with exactly three components.

Since $\kappa(G[X - \{x_1, x_2, x_3, x_4\}]) \geq \kappa(G[X]) - |\{x_1, x_2, x_3, x_4\}| \geq 16 - 4 = 12$ and $\alpha(G[X - \{x_1, x_2, x_3, x_4\}]) \leq \alpha(G) \leq 4$, $G[X - \{x_1, x_2, x_3, x_4\}]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly three components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 2.1.2. y_1 and y_2 are in different Y_i .




Figure: G has a 2-factor with exactly two components.

Since $\kappa(G[X - \{x_1, x_2, x_3, x_4\}]) \geq \kappa(G[X]) - |\{x_1, x_2, x_3, x_4\}| \geq 16 - 4 = 12$ and $\alpha(G[X - \{x_1, x_2, x_3, x_4\}]) \leq \alpha(G) \leq 4$, $G[X - \{x_1, x_2, x_3, x_4\}]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly two components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 2.1.3. $\{y_1, y_2\} \cap \{v_1, v_2\} \neq \emptyset$.




Figure: G has a 2-factor with exactly three components.


Since $\kappa(G[X - \{x_1, x_2, x_3, x_4\}]) \geq \kappa(G[X]) - |\{x_1, x_2, x_3, x_4\}| \geq 16 - 4 = 12$ and $\alpha(G[X - \{x_1, x_2, x_3, x_4\}]) \leq \alpha(G) \leq 4$, $G[X - \{x_1, x_2, x_3, x_4\}]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly three components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 2.2. $\kappa(G[Y]) = 1$.

Let v be a cut vertex of $G[Y]$. Then by $\alpha(G[Y]) = 3$, $G[Y] - v$ has at most three components.

If $G[Y] - v$ has exactly two components Y_1, Y_2 , then we can prove that both $G[Y_1]$ and $G[Y_2]$ are cliques.



Since $\kappa(G[X - \{x_1, x_2, x_3, x_4\}]) \geq \kappa(G[X]) - |\{x_1, x_2, x_3, x_4\}| \geq 16 - 4 = 12$ and $\alpha(G[X - \{x_1, x_2, x_3, x_4\}]) \leq \alpha(G) \leq 4$, $G[X - \{x_1, x_2, x_3, x_4\}]$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor with exactly two components.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Hence we assume that $G[Y] - v$ has exactly three components Y_1, Y_2 and Y_3 .

- We claim that each $G[Y_j]$ is a clique and v is adjacent to every vertex in $Y_1 \cup Y_2 \cup Y_3$.
- If $G[Y \cup N_C(Y)]$ is 2-connected, then there exists a subgraph C' of clique C such that $G[Y \cup N_C(Y) \cup C']$ is $4K_1$ -free and $|G[Y \cup N_C(Y) \cup C']| \geq r(4, 28)$, and thus $G[Y \cup N_C(Y) \cup C']$ has a 2-factor by Theorem 2.1. Also we can prove that $G[X - N_C(Y) \cup C']$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor.

Hence we assume that $\kappa(G[Y \cup N_C(Y)]) = 1$. Then there exists a vertex $x \in X \setminus (N_C(Y))$ such that x is adjacent to some vertex in $Y_1 \cup Y_2 \cup Y_3$. Let $xy_1 \in E(G)$ with $y_1 \in Y_1$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Hence we assume that $G[Y] - v$ has exactly three components Y_1, Y_2 and Y_3 .

- We claim that each $G[Y_j]$ is a clique and v is adjacent to every vertex in $Y_1 \cup Y_2 \cup Y_3$.
- If $G[Y \cup N_C(Y)]$ is 2-connected, then there exists a subgraph C' of clique C such that $G[Y \cup N_C(Y) \cup C']$ is $4K_1$ -free and $|G[Y \cup N_C(Y) \cup C']| \geq r(4, 28)$, and thus $G[Y \cup N_C(Y) \cup C']$ has a 2-factor by Theorem 2.1. Also we can prove that $G[X - N_C(Y) \cup C']$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor.

Hence we assume that $\kappa(G[Y \cup N_C(Y)]) = 1$. Then there exists a vertex $x \in X \setminus (N_C(Y))$ such that x is adjacent to some vertex in $Y_1 \cup Y_2 \cup Y_3$. Let $xy_1 \in E(G)$ with $y_1 \in Y_1$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Hence we assume that $G[Y] - v$ has exactly three components Y_1, Y_2 and Y_3 .


- We claim that each $G[Y_j]$ is a clique and v is adjacent to every vertex in $Y_1 \cup Y_2 \cup Y_3$.
- If $G[Y \cup N_C(Y)]$ is 2-connected, then there exists a subgraph C' of clique C such that $G[Y \cup N_C(Y) \cup C']$ is $4K_1$ -free and $|G[Y \cup N_C(Y) \cup C']| \geq r(4, 28)$, and thus $G[Y \cup N_C(Y) \cup C']$ has a 2-factor by Theorem 2.1. Also we can prove that $G[X - N_C(Y) \cup C']$ is hamiltonian by Chvátal-Erdős Theorem. Then G has a 2-factor.

Hence we assume that $\kappa(G[Y \cup N_C(Y)]) = 1$. Then there exists a vertex $x \in X \setminus (N_C(Y))$ such that x is adjacent to some vertex in $Y_1 \cup Y_2 \cup Y_3$. Let $xy_1 \in E(G)$ with $y_1 \in Y_1$.

Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$


Subcase 2.2.1. x has a neighbor $z_1 \in C \setminus N_C(Y)$.

We can prove that x is adjacent to every vertex in $Y_1 \cup Y_2$. Let $z_2 \in V(C) \setminus (N_C(Y) \cup \{z_1\})$. Choose a shortest path P joining z_2 and some vertex $y_3 \in Y_3$ such that the internal vertices of P are in X . Clearly, the length of P is at most three.




Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- If the length of P is two, denoted by $z_2x_1y_3$, then we can prove that x_1 is adjacent to every vertex in $Y_2 \cup Y_3$, and we can find a 2-factor of G with exactly two components.



Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- If the length of P is three, denoted by $z_2x_2x_3y_3$, then we can prove that x_2 is adjacent to some vertex in $Y_1 \cup Y_2$, and thus we can find a 2-factor of G with exactly two components.




Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

Subcase 2.2.2. $N_C(x) \subseteq N_C(Y)$.


Let x_1 be a neighbor of x in $N_C(Y)$.

- If x has a neighbor in $Y_2 \cup Y_3$, then we can prove that x is adjacent to every vertex in $Y_1 \cup Y_2 \cup Y_3$. Also we can prove that x_1 has two neighbors y_1 and y_2 in Y_1 and Y_2 , respectively, then G has a 2-factor with exactly two components.



Sketch the proof of the pair $\{K_{1,4}, P_3 \cup 2K_1\}$

- If x has no neighbor in $Y_2 \cup Y_3$, then we can prove x_1 is adjacent to every vertex in $Y_1 \cup Y_2$. Also we can prove that there exists a vertex x_2 in $N_C(Y)$ has a neighbor in Y_3 , then G has a 2-factor with exactly two components.



Thanks for your attention!