

Zdeněk Ryjáček

**On graphs with isomorphic, non-isomorphic and connected  $N_2$ -neighbourhoods**

*Časopis pro pěstování matematiky*, Vol. 112 (1987), No. 1, 66–79

Persistent URL: <http://dml.cz/dmlcz/118295>

## Terms of use:

© Institute of Mathematics AS CR, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON GRAPHS WITH ISOMORPHIC, NON-ISOMORPHIC  
AND CONNECTED  $N_2$ -NEIGHBOURHOODS

ZDENĚK RYJÁČEK, Plzeň

(Received April 18, 1984)

*Summary.* The subgraph  $N_2(u, G)$  induced by the edges  $xy$  of  $G$  for which  $\min \{\varrho(x, u), \varrho(y, u)\} = 1$  is called the neighbourhood of the second type of the vertex  $u$ . In the paper three questions are studied: existence and properties of graphs with  $N_2$ -neighbourhoods isomorphic to a given graph, existence of graphs with non-isomorphic  $N_2$ -neighbourhoods and existence and properties of graphs with connected  $N_2$ -neighbourhoods.

INTRODUCTION

Let  $G = (V(G), E(G))$  be a finite undirected graph without loops and multiple edges. The neighbourhood of an arbitrary vertex  $u \in V(G)$  (i.e. the subgraph induced on the set of vertices adjacent to  $u$ ) will be denoted by  $N_1(u)$  and called the *neighbourhood of the first type* of  $u$ . Following [2] let us denote by  $N_2(u, G)$  (or, briefly,  $N_2(u)$ ), the *neighbourhood of the second type* of  $u$ , i.e. the subgraph of  $G$  with the set of edges containing all the edges  $vw$  of  $G$  for which  $\min \{\varrho(v, u), \varrho(w, u)\} = 1$  and with the corresponding set of vertices ( $\varrho(x, y)$  denotes the distance of vertices  $x, y$ ). Then the following questions can be formulated:

1. Given a graph  $G$ , does there exist  $\tilde{G}$  such that for every vertex  $u \in V(\tilde{G})$ ,  $N_i(u)$  is isomorphic to  $G$ ? (For  $i = 1$  this is the well-known Trahtenbrot-Zykov problem, see e.g. [1], [3], [4], [5], [6].)

2. Does there exist a graph  $G$  such that for every  $u, v \in V(G)$  the neighbourhoods  $N_i(u)$  and  $N_i(v)$  are non-isomorphic? (For  $i = 1$  see [2], for 2-neighbourhoods defined as subgraphs induced on the sets of vertices at distance 2 see [7], [8].)

3. What are sufficient conditions for  $G$  to be  $N_i$ -locally connected and what are the properties of  $N_i$ -locally connected graphs? ( $G$  is said to be  $N_i$ -locally connected if for every  $v \in V(G)$  the neighbourhood  $N_i(v)$  is a connected graph. For  $i = 1$  see [9], [10].)

Investigation of these questions for  $i = 2$  is the main aim of the present paper.

## 1. $N_2$ -REALIZABLE GRAPHS

We say that a graph  $G$  is  $N_2$ -realizable if there exists a nonempty graph  $\tilde{G}$  (called an  $N_2$ -realization of  $G$ ) such that for every vertex  $u \in V(\tilde{G})$ ,  $N_2(u, \tilde{G})$  is isomorphic to  $G$ . We can assume without loss of generality that  $\tilde{G}$  is connected.

An  $N_2$ -realizable graph obviously cannot contain isolated vertices. Let us observe some other properties of  $N_2$ -realizable graphs. Denote by  $\Delta(G)$  ( $\delta(G)$ ) the maximum (minimum) degree of  $G$ .

**Theorem 1.1.** *If  $\tilde{G}$  is an  $N_2$ -realization of  $G$ , then*

$$\Delta(G) \leq \Delta(\tilde{G}) \leq \Delta(G) + 1.$$

*If moreover  $\delta(G) \geq 2$ , then*

$$\delta(\tilde{G}) \geq \delta(G) + 1.$$

**Proof.** 1. Obviously  $\Delta(G) \leq \Delta(\tilde{G})$ . Suppose that there exists  $u \in V(\tilde{G})$  such that  $d_{\tilde{G}}(u) \geq \Delta(G) + 2$  ( $d_{\tilde{G}}(u)$  denotes the degree of  $u$  in  $\tilde{G}$ ). Then  $N_2(v, \tilde{G})$  for arbitrary  $v$  adjacent to  $u$  contains a vertex of degree at least  $\Delta(G) + 1$  and therefore cannot be isomorphic to  $G$ .

2. Suppose that there exists  $u \in V(\tilde{G})$  such that  $d_{\tilde{G}}(u) \leq \delta(G)$  and consider again  $N_2(v, \tilde{G})$  for an arbitrary  $v \in V(\tilde{G})$  adjacent to  $u$ . Then the following two possibilities can occur:

a)  $N_2(v)$  does not contain  $u$ . Then  $d_{\tilde{G}}(u) = 1$  and since  $\delta(N_2(u)) \geq 2$ , necessarily  $d_{\tilde{G}}(v) \geq 3$ . Therefore  $v$  is adjacent to another vertex  $w \neq u$  and it is easily seen that  $w$  has degree 1 in  $N_2(u)$  which is a contradiction.

b)  $N_2(v)$  contains  $u$ . Then

$$\delta(\tilde{G}) \leq d_{N_2(v)}(u) = d_{\tilde{G}}(u) - 1 \leq \delta(G) - 1$$

which is again a contradiction.

**Corollary.** *An  $N_2$ -realization of a regular graph of degree  $d \geq 2$  is a regular graph of degree  $d + 1$ .*

A set  $M \subset V(G)$  is said to be a *covering set*, if every edge of  $G$  has at least one vertex in  $M$ . The minimum number of vertices in a covering set will be denoted by  $\alpha(G)$ .

**Theorem 1.2.** *If  $G$  is  $N_2$ -realizable then  $\alpha(G) \leq \Delta(G) + 1$ .*

**Proof.** If  $G = N_2(u)$ , then every edge of  $G$  has at least one vertex adjacent to  $u$  and hence the set of all vertices of  $G$  which are adjacent to  $u$  is a covering set. The proof is completed by using Theorem 1.1.

**Corollaries.** 1. If  $G$  is  $N_2$ -realizable then

$$|E(G)| \leq \Delta(G) \cdot (\Delta(G) + 1)$$

( $|M|$  denotes the number of elements of  $M$ ).

**Proof:** One vertex can cover not more  $\Delta(G)$  edges, hence  $|E(G)| \leq \alpha(G) \cdot \Delta(G)$  and we can use Theorem 1.2.

2. If  $G$  is an  $N_2$ -realizable regular graph of degree  $d$  then  $|V(G)| \leq 2(d + 1)$ .

**Proof.** Use Corollary 1 for  $\Delta(G) = d$ ,  $|E(G)| = \frac{1}{2}|V(G)| \cdot d$ .

3. a) For  $n \geq 7$  the circuit  $C_n$  is not  $N_2$ -realizable.

b) If  $G$  is a cubic (i.e. regular of degree 3) graph and  $|V(G)| \geq 9$  then  $G$  is not  $N_2$ -realizable.

c) For  $n \geq 7$  the path  $P_n$  is not  $N_2$ -realizable.

Denote by  $g(G)$  the girth of  $G$ , i.e. the length of the shortest circuit in  $G$  (if  $G$  contains no circuits, put  $g(G) = \infty$ ).

**Theorem 1.3.** Suppose  $\tilde{G}$  is an  $N_2$ -realization of  $G \neq C_3$  and  $\delta(\tilde{G}) \geq 3$ . Then  $G$  contains a path of length 3 if and only if  $g(\tilde{G}) \leq 4$ .

**Proof.** Let  $P \subset \tilde{G}$  be a path of length 3,  $P \subset N_2(u)$ . Then the vertices of  $P$  adjacent to  $u$  together with  $u$  determine in  $\tilde{G}$  a circuit of length at most 4. The converse is evident.

**Theorem 1.4.** If  $G$  is an  $N_2$ -realizable regular graph of degree  $d \geq 2$  then  $G$  is 2-connected.

**Proof.** 1. Suppose  $G$  is disconnected. For every regular graph  $G'$  of degree  $d$  we obviously have  $|V(G')| \geq d + 1$  which together with Corollary 2 of Theorem 1.2 shows that  $G$  has 2 components (each of them on  $d + 1$  vertices) and hence  $G = 2K_{d+1}$ . From  $\alpha(K_n) = n - 1$  and from Theorem 1.2 it follows that  $G$  is not  $N_2$ -realizable.

2. Suppose  $G$  has an articulation (cutvertex)  $x$ . Since each of the blocks of  $G$  has (including  $x$ ) at least  $d + 1$  vertices and  $|V(G)| \leq 2(d + 1)$ , necessarily one of the blocks of  $G$  has exactly  $d + 1$  vertices. Hence the degree-sequence of this block is

$$\underbrace{d, d, \dots, d}_d, \alpha$$

$d$ -times

for some  $\alpha < d$ , which can be easily proved to be impossible.

We shall further use the following simple assertion:

**Theorem 1.5.** Suppose  $|E(G)| \geq 1$  and let  $\tilde{G}, \tilde{\tilde{G}}$  be  $N_2$ -realizations of  $G$  such that  $\tilde{G} \subset \tilde{\tilde{G}}$ . Then  $\tilde{G} = \tilde{\tilde{G}}$ .

Proof is easy.

One can easily observe that the unique  $N_2$ -realization of the complete graph  $K_n$  for  $n > 2$  is  $K_{n+1}$ . (Here and in the sequel the term “unique” is meant up to isomorphism.) Let us consider  $N_2$ -realizability of some other classes of graphs.

**Theorem 1.6.** *The circuits  $C_3, C_5, C_6$  have a unique  $N_2$ -realization while  $C_4$  and  $C_n$  for  $n \geq 7$  are not  $N_2$ -realizable.*

**Proof.**  $n = 3$ : Let  $N_2(u) \simeq C_3$  ( $\simeq$  denotes isomorphism). We have (up to isomorphism) the following two possibilities:  $d_G(u) = 2$  or  $d_G(u) = 3$ . In the first case we obtain an  $N_2$ -realization of  $C_3$  isomorphic to  $K_4$ , in the second case considering  $N_2(v)$  of any vertex  $v$  adjacent to  $u$  we are led again to an  $N_2$ -realization isomorphic to  $K_4$ .

$n = 4$ : Let  $N_2(u) \simeq C_4$ . Then some two non-adjacent vertices  $v_1, v_2$  of  $C_4$  must be joined with  $u$  by an edge, which implies that  $v_2$  has degree 3 in  $N_2(v_1)$  – a contradiction.

$n = 5$ : If  $N_2(u) \simeq C_5$  then there necessarily exist three vertices  $v_1, v_2, v_3$  on  $C_5$  such that (say)  $v_1$  is not adjacent to  $v_2$  and  $v_3$  but  $v_2$  is adjacent to  $v_3$  and all of them are adjacent to  $u$ . Considering  $N_2(v_1)$  and using Theorem 1.5 we obtain the only possible  $N_2$ -realization to be  $C_3 \times P_1$ , i.e. the graph of the trigonal prism.

$n = 6$ : Similarly as in the preceding case it can be proved that the only  $N_2$ -realization of  $C_6$  is the graph of the 3-dimensional cube.

For  $n \geq 7$  see Corollary 3a of Theorem 1.2.

A vertex  $u \in V(G)$  is said to be *universal* if it is adjacent to all other vertices of  $G$ .

**Theorem 1.7.** *If  $G$  has exactly one universal vertex and  $|V(G)| = n \geq 4$ , then one of the following possibilities occurs:*

- a)  $G \simeq K_{1, n-1}$  and  $G$  is uniquely  $N_2$ -realizable;
- b)  $n$  is odd,  $G \simeq K_{\underbrace{2, 2, \dots, 2}_1, 1}$  and  $G$  has the unique

$$N_2\text{-realization } \tilde{G} \simeq K_{\underbrace{2, 2, \dots, 2}_{\frac{1}{2}(n-1) \text{ times}}};$$

- c)  $G$  is not  $N_2$ -realizable.

**Proof.** Suppose that  $N_2(u_0) \simeq G$  has  $n$  vertices  $u_1, \dots, u_n$ ,  $u_1$  is universal in  $N_2(u_0)$  and  $\tilde{G}$  is an  $N_2$ -realization of  $G$ .

Case 1. Suppose  $u_0, u_1$  are adjacent in  $\tilde{G}$ . Then the neighbourhood  $N_2(u_1)$  must have a universal vertex and without loss of generality we may assume that it is  $u_0$ . If there exists a vertex  $u_k$  ( $k \neq 0, 1$ ) which is adjacent to both  $u_0$  and  $u_1$  then an easy

consideration shows that both  $u_0$  and  $u_1$  are universal in  $N_2(u_k)$  which is a contradiction. Hence no  $u_k$  is adjacent to both  $u_0$  and  $u_1$  and by considering  $N_2(u_1)$  and using Theorem 1.5 it is seen that the only possible  $\tilde{G}$  is the “double-star”, i.e. the tree consisting of the edge  $u_0u_1$ ,  $n - 1$  edges  $u_ku_1$  for  $2 \leq k \leq n$  and  $n - 1$  other edges adjacent to  $u_0$ ; the resulting graph is an  $N_2$ -realization of the star  $K_{1,n-1}$ .

Case 2. If  $u_0, u_1$  are not adjacent in  $\tilde{G}$  then the universality of  $u_1$  in  $N_2(u_0)$  implies that  $u_0$  is adjacent to all  $u_i$  for  $i = 2, \dots, n$ . Now the neighbourhood  $N_2(u_i)$  for every  $i = 0, 1, \dots, n$  has exactly  $n$  vertices and hence  $\tilde{G}$  cannot have any other vertices. We shall prove by induction the following assertion:

**Lemma.** *Let  $l$  be an integer such that  $1 \leq l \leq \frac{1}{2}(n - 1)$ . If each of the graphs  $N_2(u_i)$ ,  $i = 0, \dots, 2l - 1$  contains exactly one universal vertex then all pairs of vertices  $u_i, u_j$  for  $0 \leq i \leq 2l - 1$  are adjacent in  $G$  except the pairs  $u_{2k}, u_{2k+1}$  for  $k = 0, 1, \dots, l - 1$ .*

**Proof.** For  $l = 1$  the lemma holds evidently. Suppose that  $l \leq \frac{1}{2}(n - 1)$  and the assertion of our lemma is true for  $l - 1$  – therefore the pairs of vertices  $u_{2k}, u_{2k+1}$  are not adjacent for  $k = 0, 1, \dots, l - 2$ . This implies that none of the vertices  $u_i$  for  $i < 2l - 2$  can be universal in  $N_2(u_{2l-2})$ ; hence this universal vertex must be one of  $u_i$  for  $2l - 1 \leq i \leq n$  and we may assume without loss of generality that it is  $u_{2l-1}$ . Hence  $u_{2l-1}$  is adjacent in  $\tilde{G}$  to all  $u_j$  for  $2l \leq j \leq n$  and therefore the vertices  $u_{2l-2}, u_{2l-1}$  cannot be adjacent in  $\tilde{G}$  (since in the other case  $u_{2l-1}$  would be another universal vertex in  $N_2(u_0)$ ). This implies that all the pairs  $u_{2l-2}, u_j$  for  $2l \leq j \leq n$  are adjacent in  $\tilde{G}$  and the lemma is proved.

Case 2a.  $n$  is odd. Using our lemma for  $l = \frac{1}{2}(n - 1)$  and observing that the vertices  $u_{n-1}, u_n$  cannot be adjacent in  $\tilde{G}$  (since otherwise both  $u_{n-1}$  and  $u_1$  would be universal in  $N_2(u_0)$ ) it is proved that the only possibility is  $G \simeq \underbrace{K_{2,2, \dots, 2}}_{\frac{1}{2}(n+1) \text{ times}}$ .

$$\frac{1}{2}(n+1) \text{ times}$$

Case 2b.  $n$  is even. Then using the lemma for  $l = \frac{1}{2}(n - 2)$  and considering  $N_2(u_{n-2})$  we conclude that one of  $u_{n-1}, u_n$  (say  $u_{n-1}$ ) must be universal in  $N_2(u_{n-2})$ . Then  $u_{n-1}, u_n$  and one of the pairs of vertices  $u_{n-2}, u_{n-1}$  and  $u_{n-2}, u_n$  must be adjacent. But in the first case  $u_{n-1}$  and in the other case  $u_n$  is another universal vertex in  $N_2(u_0)$ . This contradiction proves the non-existence of an  $N_2$ -realization.

**Corollary.** *The wheels  $W_3$  and  $W_4$  are uniquely  $N_2$ -realizable while  $W_n$  for  $n \geq 5$  is not  $N_2$ -realizable (wheel  $W_n$  is  $C_n$  together with an additional univesal vertex).*

**Proof.**  $\tilde{W}_3 \simeq K_5$  since  $W_3 \simeq K_4$ ,  $\tilde{W}_4 \simeq K_{2,2,2}$  since  $W_4 \simeq K_{2,2,1}$ , for  $n \geq 5$  use Theorem 1.7.

**Theorem 1.8.** *Let  $G$  be a disjoint union of stars, i.e.*

$$G = \bigcup_{i=1}^n K_{k_i,1}, \quad k_i \geq 2, \quad i = 1, \dots, n, \quad n \geq 2.$$

*Then  $G$  is  $N_2$ -realizable if and only if  $k_1 = k_2 = \dots = k_n = n - 1$  and in this case  $G$  has infinitely many non-isomorphic  $N_2$ -realizations.*

**Proof.** Suppose  $G$  is  $N_2$ -realizable. First observe that if  $\tilde{G}$  is an  $N_2$ -realization of  $G$  then an arbitrary vertex  $u \in V(\tilde{G})$  is adjacent in  $\tilde{G}$  to all centers of components of  $G$  and to no other vertices: if some end-vertex  $v$  of  $G$  were adjacent to  $u$  in  $\tilde{G}$  then its neighbourhood  $N_2(v)$  should contain a path of length 3 which is a contradiction. Hence  $\tilde{G}$  is a regular graph of degree  $n$  and therefore necessarily  $k_1 = k_2 = \dots = k_n = n - 1$ .

Conversely, suppose  $k_1 = k_2 = \dots = k_n = n - 1$ . Then  $G = nK_{n-1,1}$  and according to Theorem 1.3,  $G$  is  $N_2$ -realized by an arbitrary regular graph  $\tilde{G}$  of degree  $n$  such that  $g(\tilde{G}) \geq 5$ . Existence of an infinite family of such graphs is proved in [12], Chapter III, Theorem 1.4'.

Denote by  $P_k$ ,  $k \geq 1$ , the path of length  $k$ , i.e. with  $k$  edges and  $k + 1$  vertices.

**Theorem 1.9.** *Let  $G$  be a disjoint union of paths, i.e.  $G = \bigcup_{i=1}^n P_{k_i}$ ,  $k_i \geq 1$ ,  $i = 1, \dots, n$ ,  $n \geq 1$ . Then  $G$  is  $N_2$ -realizable only in the following cases:*

$n$ (number of paths)	$k_i$ ( $i = 1, \dots, n$ ) (lengths of paths)	number of non-isomorphic $N_2$ -realizations
1	1	2
	2	2
	3	1
	6	$\infty$
2	1, 1	$\infty$
	2, 3	$\infty$
	2, 4	$\infty$
3	2, 2, 2	$\infty$

**Proof.** If  $\tilde{G}$   $N_2$ -realizes  $G$  then according to Theorem 1.2 necessarily  $\alpha(G) \leq 3$ . Hence  $n \leq 3$  and it remains to consider the following possibilities: for  $n = 1$ :  $k = 1, 2, 3, 4, 5, 6$ ; for  $n = 2$ :  $k_i = 1, 1; 1, 2; 1, 3; 1, 4; 2, 2; 2, 3; 2, 4$ ; for  $n = 3$ :  $k_i = 1, 1, 1; 1, 1, 2; 1, 2, 2; 2, 2, 2$ .

Case  $n = 1$ . Non-realizability of  $P_4$  and  $P_5$  is proved and examples of  $N_2$ -realizations of  $P_1, P_2, P_3$  and  $P_6$  are given in [2]. It remains to prove the assertion concerning the number of  $N_2$ -realizations.

a) Let  $N_2(u) \simeq P_1$ . Then  $u$  is adjacent either to one of the vertices of  $P_1$  or to both of them. In virtue of Theorem 1.5 the first case yields  $C_3$  and the second case yields  $P_3$  as the only possible  $N_2$ -realizations.

b) Let  $N_2(u) \simeq P_2$ , let  $v_1, v_2, v_3$  be the three vertices of  $P_2$ . We have (up to isomorphism) the following four possibilities:  $u$  is adjacent to  $v_2$ ;  $u$  is adjacent to  $v_1$  and  $v_2$ ;  $u$  is adjacent to  $v_1$  and  $v_3$ ;  $u$  is adjacent to  $v_1, v_2$  and  $v_3$ . In the first case considering  $N_2(v_2)$  we obtain the first  $N_2$ -realization of  $P_2$  which is a tree on 6 vertices with exactly 2 of them of degree 3 while in the third case we obtain  $C_4$  as the second possible  $N_2$ -realization of  $P_2$ . The second and fourth cases imply a contradiction.

c) Let  $N_2(u) \simeq P_3$ . In a similar manner as in the preceding case it can be proved that the  $N_2$ -realization which is shown in [2] (i.e. the circuit  $C_5$  with one diagonal edge) is the only one.

d) In [2] it is shown that  $P_6$  is  $N_2$ -realized by the graph of the  $m$ -gonal prism  $C_m \times P_1$  for arbitrary  $m \geq 5$ .

Case  $n = 2$ . a)  $C_m$   $N_2$ -realizes  $2P_1$  for an arbitrary  $m \geq 5$ .

b) Suppose  $N_2(u) \simeq P_1 \cup P_2$ ,  $V(P_1) = \{v_1, v_2\}$ ,  $V(P_2) = \{w_1, w_2, w_3\}$ . According to Theorem 1.1  $d_{\mathcal{G}}(u) \leq 3$  and hence we obtain the following three possibilities:

- $u$  is adjacent to  $w_2$  and one of  $v_i$ 's (say  $v_1$ );
- $u$  is adjacent to  $w_1, w_3$  and one of  $v_i$ 's (say  $v_1$ );
- $u$  is adjacent to  $w_2, v_1$  and  $v_2$ .

The last two cases immediately imply a contradiction while in the first case the condition  $N_2(w_2) \simeq P_1 \cup P_2$  implies that either one of the vertices  $w_1, w_3$  must have degree 1 or they are joined by another path  $P_2$ . In both of these cases considering  $N_2(w_1)$  we obtain a contradiction.

c) Let  $N_2(u) \simeq P_1 \cup P_3$ . Then necessarily  $d_{\mathcal{G}}(u) = 3$ . Since  $N_2(u) \supset P_1$ , the vertex  $u$  is adjacent to some vertex  $v$  of degree 2 in  $\mathcal{G}$  and hence  $N_2(v)$  cannot be isomorphic to  $P_1 \cup P_3$ .

d) Non-realizability of  $P_1 \cup P_4$  can be proved similarly.

e) Non-realizability of  $2P_2 \simeq 2K_{2,1}$  follows from Theorem 1.8.

f) An  $N_2$ -realization of the graph  $P_2 \cup P_3$  can be constructed by using an arbitrary connected regular graph of degree 3 and replacing each of its vertices by  $C_3$ .

g) An  $N_2$ -realization of the graph  $P_2 \cup P_4$  can be constructed in a similar manner as in the above case by using a connected regular graph of degree 4 and the circuit  $C_4$ .

Case  $n = 3$ . a) If  $N_2(u)$  is a graph with 3 components and one of them is  $P_1$  then  $u$  is adjacent to some vertex  $v$  such that  $d_{\mathcal{G}}(v) = 2$  and hence  $N_2(v)$  cannot be a graph with 3 components. Hence the graphs  $3P_1, 2P_1 \cup P_2$  and  $P_1 \cup 2P_2$  are not  $N_2$ -realizable.



b)  $3P_2 \simeq 3K_{2,1}$  has infinitely many  $N_2$ -realizations according to Theorem 1.8.

**Theorem 1.10.** *The complete bipartite graph  $K_{m,n}$  is  $N_2$ -realizable if and only if either  $\min\{m, n\} = 1$  or  $|m - n| = 1$ . The graphs  $K_{1,1}$  and  $K_{1,2} \simeq K_{2,1}$  have exactly two non-isomorphic  $N_2$ -realizations while in the other cases the  $N_2$ -realization of  $K_{m,n}$  is unique.*

**Proof.** The assertion concerning  $K_{1,1} \simeq P_1$  and  $K_{1,2} \simeq K_{2,1} \simeq P_2$  follows from Theorem 1.9 while the assertion concerning  $K_{1,n} \simeq K_{n,1}$  for  $n \geq 3$  follows from Theorem 1.7.

Let  $\tilde{G}$  be an  $N_2$ -realization of  $G = K_{m,n}$ ,  $u_0 \in V(\tilde{G})$ ,  $N_2(u_0) \simeq K_{m,n}$ ,  $m \geq 2$ ,  $n \geq 2$ . Let  $A = \{a_1, \dots, a_m\}$ ,  $B = \{b_1, \dots, b_n\}$  be the two classes of vertices of  $K_{m,n}$ . Then  $u_0$  is adjacent either to all  $a_i$ 's or to all  $b_j$ 's since otherwise for a pair of vertices  $a_{i_0}, b_{j_0}$  such that none of them is adjacent to  $u_0$  the edge  $a_{i_0}b_{j_0}$  would not be in  $N_2(u_0)$ . Further,  $u_0$  is adjacent either to all  $a_i$ 's and no  $b_j$ 's or to all  $b_j$ 's and no  $a_i$ 's since in the first case for  $b_{j_0}$  adjacent to  $u_0$  the neighbourhood  $N_2(a_1)$  would contain the circuit of length 3 with vertices  $a_2, b_{j_0}, u_0$ ; the second case is similar. Consequently, in the first case  $a \in A \Rightarrow N_2(a) \simeq K_{m-1, n+1}$ ,  $b \in B \Rightarrow N_2(b) \simeq N_2(u_0) \simeq K_{m,n}$  and hence  $m - n = 1$  and  $\tilde{G} \simeq K_{m,m}$ ; in the second case  $a \in A \Rightarrow N_2(a) \simeq N_2(u_0) \simeq K_{m,n}$ ,  $b \in B \Rightarrow N_2(b) \simeq K_{m+1, n-1}$  and hence  $n - m = 1$  and  $\tilde{G} \simeq K_{n,n}$ .

**Theorem 1.11.** *The only  $N_2$ -realizable cubic (i.e. regular of degree 3) graphs are the tetrahedron  $K_4$ , the trigonal prism  $C_3 \times P_1$  and the 3-dimensional cube  $Q_3$ , and each of them has a unique  $N_2$ -realization.*

**Proof.** The only cubic graph with four vertices is the uniquely  $N_2$ -realizable tetrahedron  $K_4$ . For  $|V(G)| = 6$  there exist 2 non-isomorphic cubic graphs, namely




Fig. 1

$K_{3,3}$  and the trigonal prism  $C_3 \times P_1$ .  $K_{3,3}$  is not  $N_2$ -realizable according to Theorem 1.10. Suppose  $N_2(u) \simeq C_3 \times P_1$ ,  $u_{i,j}$  ( $i = 1, 2, 3, j = 1, 2$ ) being its vertices.

Necessarily  $d_{\tilde{G}}(u) = 4$  and hence the only (up to isomorphism) possibility is that  $u_{1,1}, u_{2,1}, u_{2,2}$  and  $u_{3,2}$  are adjacent to  $u$  (these vertices must form a covering set). The condition  $N_2(u_{1,1}) \simeq C_3 \times P_1$  then implies that the vertices  $u_{3,1}$  and  $u_{1,2}$  are adjacent in  $\tilde{G}$  and hence we have obtained the unique  $N_2$ -realization which is shown in Fig. 1.

If  $|V(G)| = 8$  then  $\alpha(G) = 4$  according to Theorem 1.2 and hence  $G$  is necessarily bipartite. The only bipartite cubic graph with 8 vertices in the 3-dimensional cube the  $N_2$ -realization of which is shown in Fig. 2. The proof of uniqueness is similar to the preceding case.




Fig. 2

For  $|V(G)| > 8$  see Corollary 3b of Theorem 1.2.

### Corollaries.

1. *The only  $N_2$ -realizable cube  $Q_n$  is the 3-dimensional one.*

*Proof.* For  $Q_2 \simeq C_4$  and  $Q_3$  see Theorems 1.6 and 1.11.  $Q_n$  is not  $N_2$ -realizable for  $n \geq 4$  according to Corollary 2 of Theorem 1.2 since  $Q_n$  is regular of degree  $n$  and  $|V(Q_n)| = 2^n > 2(n + 1)$ .

2. *The only  $N_2$ -realizable graphs of Platonic bodies are the tetrahedron and the cube, and their  $N_2$ -realizations are unique.*

*Proof.*  $N_2$ -realizability of the tetrahedron and the cube and non-realizability of the dodecahedron is established by the preceding theorem. The icosahedron is not  $N_2$ -realizable since it has no covering with at most 6 vertices. Suppose  $G$  is the graph of the octahedron,  $\tilde{G}$  its  $N_2$ -realization,  $u \in V(\tilde{G})$ ,  $N_2(u) \simeq G$ . Necessarily  $d_{\tilde{G}}(u) = 5$ ; hence we may denote by  $u_1$  the vertex of  $G$  which is not adjacent to  $u$ , by  $u_6$  the only vertex of  $G$  which is not adjacent to  $u_1$ , and by  $u_2, u_3, u_4, u_5$  the other vertices of  $G$ .  $\tilde{G}$  is regular of degree 5 and hence  $u_1$  is necessarily adjacent in  $\tilde{G}$  either to  $u_6$  or to

another vertex  $v$ , but it can be shown that both of these possibilities lead to a contradiction.

## 2. GRAPHS WITH NON-ISOMORPHIC $N_2$ -NEIGHBOURHOODS

Following [2] let us denote by  $\mathfrak{G}_2$  the class of graphs with the following property: for every pair of vertices  $u, v$  of  $G$  the neighbourhoods  $N_2(u)$  and  $N_2(v)$  are not isomorphic.

**Theorem 2.1.** *Let  $n$  be an integer. Then there exists a connected graph  $G_n$  on  $n$  vertices belonging to  $\mathfrak{G}_2$  if and only if  $n \geq 7$ .*

We shall first prove some auxiliary assertions.

**Lemma 1.** *Let  $n \geq 7$ ,  $G_n \in \mathfrak{G}_2$ , suppose that  $G_n$  is connected, none of the vertices  $u_1, \dots, u_n$  of  $G_n$  is universal and the only vertex which is adjacent to  $u_n$  is  $u_{n-2}$ . Let us construct a graph  $G_{n+1}$  on  $n + 1$  vertices from  $G_n$  by adding a vertex  $u_{n+1}$  and making it universal in  $G_{n+1}$ . Then  $G_{n+1} \in \mathfrak{G}_2$ ,  $G_{n+1}$  is connected and  $u_n$  is adjacent only to  $u_{n-2}$  and  $u_{n+1}$ .*

*Proof.* Suppose that  $f: N_2(u_\alpha, G_{n+1}) \rightarrow N_2(u_\beta, G_{n+1})$  is an isomorphism. Without loss of generality we may assume that  $\alpha \neq n + 1$  and hence  $u_{n+1} \in V(N_2(u_\alpha, G_{n+1}))$ . If  $f(u_{n+1}) = u_{n+1}$  then the partial mapping  $f|_{V(G_n)}$  is an isomorphism  $N_2(u_\alpha, G_n)$  onto  $N_2(u_\beta, G_n)$ . Hence  $f(u_{n+1}) = u_\gamma$ ,  $\gamma \leq n$  and  $u_\gamma$  is universal in  $N_2(u_\beta, G_{n+1})$ . If  $\beta = n + 1$  then  $N_2(u_\beta, G_{n+1}) = G_n$  and  $u_\gamma$  is universal in  $G_n$ . Hence  $\beta \leq n$  and therefore  $u_{n+1}$  is the second universal vertex in  $N_2(u_\beta, G_{n+1})$ . Interchanging these two universal vertices we obtain an isomorphism  $f_1: N_2(u_\alpha, G_{n+1}) \rightarrow N_2(u_\beta, G_{n+1})$  such that  $f_1(u_{n+1}) = u_{n+1}$ , which is a contradiction.

**Lemma 2.** *Let  $n \geq 7$ ,  $G_n \in \mathfrak{G}_2$ ,  $V(G_n) = \{u_1, \dots, u_n\}$ , suppose that  $u_n$  is universal in  $G_n$ , the only vertex of degree 1 in  $N_2(u_n, G_n)$  is  $u_{n-1}$  and  $u_{n-1}$  is adjacent only to  $u_{n-3}$  and  $u_n$ . Let us construct a graph  $G_{n+1}$  on  $n + 1$  vertices from  $G_n$  by adding a vertex  $u_{n+1}$  and joining it to  $u_{n-1}$  by an edge. Then  $G_{n+1} \in \mathfrak{G}_2$ ,  $G_{n+1}$  is connected and has no universal vertex.*

*Proof.* The vertex  $u_n$  is universal in  $G_n$  and hence all vertices of  $G_n$  have (by assumption, non-isomorphic)  $N_2$ -neighbourhoods on  $n - 1$  vertices. The only vertices  $u_i$  of  $G_{n+1}$  for which  $N_2(u_i, G_{n+1}) \neq N_2(u_i, G_n)$  are evidently  $u_{n-3}$  and  $u_n$  (and, of course,  $u_{n+1}$ ).  $N_2(u_{n+1}, G_{n+1})$  has 3 vertices while both  $N_2(u_{n-3}, G_{n+1})$  and  $N_2(u_n, G_{n+1})$  have  $n$  vertices. Suppose that there exists an isomorphism  $f: N_2(u_n, G_{n+1}) \rightarrow N_2(u_{n-3}, G_{n+1})$ . By assumption, the only vertex of degree 1 in both  $N_2(u_n, G_{n+1})$  and  $N_2(u_{n-3}, G_{n+1})$  is  $u_{n+1}$ . Hence the partial mapping  $f|_{V(G_n)}$  is an isomorphism of  $N_2(u_n, G_n)$  onto  $N_2(u_{n-3}, G_n)$ , which is a contradiction.

Proof of Theorem 2.1. The non-existence of  $G_n \in \mathfrak{G}_2$  for  $n \leq 6$  can be easily verified by listing all such graphs (see e.g. [13]). For  $n \geq 7$  let us construct a graph  $G_n$  using the following construction:

– for  $n = 7$  see Fig. 3;




Fig. 3

– having obtained  $G_n$ , construct  $G_{n+1}$  using Lemma 1 if  $n$  is odd and Lemma 2 if  $n$  is even.

Then  $G_{n+1}$  is connected and  $G_{n+1} \in \mathfrak{G}_2$ .

**Theorem 2.2.** Let  $n, k$  be integers,  $k \geq 1$ ,  $n \geq k^2 + 5k + 1$ . Then there exists a graph  $G \in \mathfrak{G}_2$  with  $n$  vertices and  $k$  components.

Proof. Let us define a graph  $G$  using the graphs  $G_n$  which are described in the proof of Theorem 2.1:

- the first component of  $G$  is  $G_7$ ,
- the  $i$ -th component of  $G$  is  $G_{2i+4}$ ,  $i = 2, \dots, k$ .

Then every component of  $G$  belongs to  $\mathfrak{G}_2$  and since for every pair of vertices  $u_1, u_2$  which belong to different components of  $G$  their  $N_2$ -neighbourhoods have different numbers of vertices, necessarily  $G \in \mathfrak{G}_2$ . Further,  $n = 7 + \sum_{i=2}^k (2i + 4) = k^2 + 5k + 1$  and hence for  $n = k^2 + 5k + 1$  the theorem is proved.

For  $n > k^2 + 5k + 1$  take the same graph  $G$  with the only difference in the  $k$ -th component: if we denote  $a = n - (k^2 + 5k + 1)$  then it is constructed as  $G_{2k+4+a}$  if  $a$  is even and as a graph which can be obtained from  $G_{2k+3+a}$  by adding a new vertex and joining it to the only universal vertex of  $G_{2k+3+a}$  if  $a$  is odd.

### 3. $N_2$ -LOCALLY CONNECTED GRAPHS

**Theorem 3.1.** Let  $G$  be a connected  $N_2$ -locally connected graph, suppose that  $G$

contains a path of length 4. Denote by  $G'$  the graph which is obtained from  $G$  by deleting all vertices of degree 1 together with their edges. Then every edge of  $G'$  is contained in some circuit of length  $m \leq 4$  and  $G'$  is 2-connected.

*Proof.* Let  $h$  be an edge of  $G'$ . Each of its vertices is adjacent to another edge – denote them by  $h_1, h_2$ . If  $h_1, h_2$  have a common vertex then  $h$  is contained in a triangle  $h, h_1, h_2$ . Suppose that  $h_1, h_2$  have no common vertex and that in  $G$  there is no circuit of length  $m \leq 4$  containing  $h$ . Then the existence of path of length 4 in  $G$  and the connectedness of  $G$  yield that in  $G$  there exists a path of length 4 such that if  $u_0, u_1, u_2, u_3, u_4$  are its vertices then  $h = u_1u_2$ . The neighbourhood  $N_2(u_2, G)$  then contains the edges  $u_0u_1$  and  $u_3u_4$ . Suppose that in  $G$  there is no circuit of length  $m \leq 4$  containing  $h$ . Hence if a vertex  $v$  is adjacent to  $u_1$  and  $w$  is adjacent to  $u_2$  then  $v$  cannot be adjacent to  $w$  and therefore the edges  $u_0u_1$  and  $u_3u_4$  are in different components of  $N_2(u_2, G)$ .

Let  $u$  be an articulation of  $G'$ . Then  $u$  is an articulation of  $G$  and such edges  $h_1, h_2$  can be found that  $h_1, h_2$  are in different blocks of  $G$  and none of them is adjacent to  $u$  (since otherwise  $u$  would not be an articulation of  $G'$ ). But then  $N_2(u, G)$  is disconnected, which is a contradiction.

Obviously, every  $N_1$ -locally connected graph  $G$  is  $N_2$ -locally connected and hence the assertions which are proved in [9], [10] can be used to obtain sufficient conditions for  $G$  to be  $N_2$ -locally connected. Nevertheless, some of them can be replaced by weaker ones.

**Theorem 3.2.** *Every graph which contains no path of length 4 is  $N_2$ -locally connected.*

*Proof* is easy.

**Theorem 3.3.** *Let  $G$  be a graph such that every pair  $u, v$  of non-adjacent vertices satisfies the inequality*

$$d_G(u) + d_G(v) \geq |V(G)|.$$

*Then  $G$  is  $N_2$ -locally connected.*

*Proof.* Let  $u_0 \in V(G)$  and suppose that  $N_2(u_0, G)$  is disconnected. Choose vertices  $u_1, u_2$  in different components of  $N_2(u_0)$  so that they are adjacent to  $u_0$ . Each of the vertices  $u_1, u_2$  is adjacent to  $d_G(u_i) - 1$  vertices (excluding  $u_0$ ) and these vertices are necessarily different. Hence

$$|V(G)| \geq (d_G(u_1) - 1) + (d_G(u_2) - 1) + 3$$

which implies

$$d_G(u_1) + d_G(u_2) \leq |V(G)| - 1,$$

a contradiction.

**Example.** The graph  $G$  which can be obtained by taking two disjoint copies of  $K_n$ ,  $n \geq 2$ , and joining their vertices with an additional universal vertex  $u$ , is not  $N_2$ -locally connected and every pair  $x, y$  its of vertices such that  $x \neq u$  and  $y \neq u$  satisfies  $d_G(x) + d_G(y) = 2n < 2n + 1 = |V(G)|$ . Hence Theorem 3.3 is the best possible.

**Corollary.** *If  $\delta(u) \geq \frac{1}{2}|V(G)|$  then  $G$  is  $N_2$ -locally connected.*

**Theorem 3.4.** *Let  $G$  be a graph without triangles and such that*

$$\sum_{u \in V(P)} d_G(u) \geq |V(G)| + 2$$

*for every path  $P \subset G$  of length 2. Then  $G$  is  $N_2$ -locally connected.*

**Proof.** Let  $u_0, u_1, u_2$  be the same as in the proof of Theorem 3.3. Then  $u_0$  is adjacent to  $d_G(u_0)$  vertices and each of the vertices  $u_1, u_2$  is adjacent to another  $d_G(u_i) - 1$  vertices. These vertices are different since  $N_2(u_0, G)$  is disconnected and  $G$  has no triangles. Hence

$$|V(G)| \geq d_G(u_0) + d_G(u_1) - 1 + d_G(u_2) - 1 + 1$$

which yields

$$\sum_{i=0}^2 d_G(u_i) \leq |V(G)| + 1,$$

a contradiction.

**Corollary.** *Suppose that  $G$  is a graph without triangles for which one of the following conditions is fulfilled:*

a) *for every pair of vertices  $u, v$ ,*

$$d_G(u) + d_G(v) \geq \frac{2}{3}(|V(G)| + 2);$$

b)

$$\delta(G) \geq \frac{1}{3}(|V(G)| + 2).$$

*Then  $G$  is  $N_2$ -locally connected.*

**Acknowledgment.** The author is indebted to J. Sedláček, for his helpful suggestions.

#### References

- [1] A. A. Zykov: Problem 30, Theory of graphs and its applications. Proc. Symp. Smolenice 1963 (M. Fiedler, ed.), Prague 1964, 164–165.
- [2] J. Sedláček: Local properties of graphs. Čas. pěst. mat. 106 (1981), 290–298.

- [3] *V. K. Bulitko*: On graphs with given vertex-neighbourhoods. *Trudy mat. inst. im. Steklova* 133 (1973), 78–94.
- [4] *P. Hell*: Graphs with given neighbourhoods I. *Problèmes Combinatoires et Théorie des Graphes (Colloq. Orsay 1976)*, C.N.R.S., Paris 1978, 219–223.
- [5] *J. I. Hall*: Locally Petersen graphs. *J. Graph Theory* 4 (1980), 173–187.
- [6] *A. Vinc.*: Locally homogeneous graphs from groups. *J. Graph Theory* 5 (1981), 417–422.
- [7] *H. Bielak*: On graphs with non-isomorphic 2-neighbourhoods. *Čas. pěst. mat.* 108 (1983), 294–298.
- [8] *H. Bielak, E. Soczewińska*: Some remarks about digraphs with non-isomorphic 1- or 2-neighbourhoods. *Čas. pěst. mat.* 108 (1983), 299–304.
- [9] *G. Chartrand, R. E. Pipert*: Locally connected graphs. *Čas. pěst. mat.* 99 (1974), 158–163.
- [10] *D. W. VanderJagt*: Sufficient conditions for locally connected graphs. *Čas. pěst. mat.* 99 (1974), 400–404.
- [11] *O. Ore*: *Theory of graphs*. AMS, Providence, R. I. 1962.
- [12] *B. Bollobás*: *Extremal graph theory*. Academic Press 1978.
- [13] *F. Harary*: *Graph theory*. Addison-Wesley, Reading, Mass. 1969.

### Souhrn

#### O GRAFECH S ISOMORFNÍMI, NEISOMORFNÍMI A SOUVISLÝMI $N_2$ -OKOLÍMI

ZDENĚK RYJÁČEK

Podgraf  $N_2(u, G)$  grafu  $G$  indukovaný množinou hran  $xy$  grafu  $G$ , pro něž  $\min\{\varrho(x, u), \varrho(y, u)\} = 1$ , se nazývá okolí 2. druhu uzlu  $u$ . V článku jsou vyšetřovány tři otázky: existence a vlastnosti grafů, v nichž  $N_2$ -okolí každého uzlu je isomorfní z daným grafem, existence grafů s neisomorfními  $N_2$ -okolími uzlů a existence a vlastnosti grafů, v nichž  $N_2$ -okolí všech uzlů jsou souvislá.

### Резюме

#### О ГРАФАХ С ИЗОМОРФНЫМИ, НЕИЗОМОРФНЫМИ И СВЯЗНЫМИ $N_2$ -ОКРУЖЕНИЯМИ

ZDENĚK RYJÁČEK

Подграф  $N_2(u, G)$ , порожденный такими ребрами  $xy$  графа  $G$ , для которых  $\min\{\varrho(x, u), \varrho(y, u)\} = 1$ , называется окружением второго типа вершины  $u$ . В настоящей статье рассмотрены следующие три вопроса: существование и свойства графов,  $N_2$  — окружения вершин которых изоморфны заданному графу, существование графов,  $N_2$  — окружения вершин которых неизоморфны и существование и свойства графов,  $N_2$  — окружения вершин которых являются связными.

*Author's address*: 306 14 Plzeň, Nejedlého sady 14 (Katedra matematiky VŠSE).